




Speleological Association of Slovenia
and
Karst Research Institute ZRC SAZU

2nd INTERNATIONAL KARSTOLOGICAL SCHOOL

Classical Karst



GUIDE-BOOKLET FOR THE EXCURSIONS

Postojna, June 27 - 30, 1994

2nd INTERNATIONAL KARSTOLOGICAL SCHOOL
Classical Karst

Contents

INTRODUCTION

PROGRAM

ABSTRACTS OF PAPERS

MATERIALS FOR THE FIELD-WORK

RELATED REFERENCES

Supported by: Ministry of Science and Technology and
the Postojna Commune

<i>Publisher:</i>	Inštitut za raziskovanje krasa ZRC SAZU Tilov trg 2 66 230 Postojna
<i>Composed by:</i>	A. Kranjc, A. Mihevc
<i>Translated by:</i>	M. Kranjc

INTRODUCTION

The 1st International Karstological School last year was dedicated to Kras, to the landscape which gave the name to term "kras" and to the related science - karstology. The purpose of our international summer schools is to acquaint the participants with the entire Slovene classical karst. For this reason we selected as a basic topic of this year the karst poljes.

Karst polje is one of the rudimentary and at the same time the most noticeable karst morphological feature of Dinaric karst, consequently of the Slovene classical karst. As in many karst types the karst poljes did not develop they are extremely important for cognition of the classical karst and its morphological, and in particular hydrological genesis.

The expression "polje" itself is the word that arises from the Slavic languages and has its meaning in today's slovene language too. Moreover polje is the morphological feature where human activity and karst nature are the most tightly associated. From old the karst poljes with the flat surfaces, covered by the sediments, attracted the farmers and at the same time discourage them due to threatening and unpredictable floods, to settle or to cultivate them more intensively. An important role of this year's karstological school is to throw light upon this relation and at the same time the relation man - nature is an important element entitling the incorporation of this international school to UNESCO.

Andrej Kranjc

PROGRAM OF THE 2nd INTERNATIONAL KARSTOLOGICAL SCHOOL

Postojna, June 27 - 30, 1994

MONDAY, June 27

- 9.00 opening
10.00 coffee break
Lectures

10.30 Ivan Gams, TYPES OF POLJES IN SLOVENIA AND
THEIR INFLUENCE ON LAND USE
11.00 France Šušteršič, POLJES AND CAVES OF NOTRANJSKA
11.30 Jože Čar, STRUCTURE OF IDRJA FAULT ZONE ON
PLANINSKO POLJE

13.00 Field work - Planinsko polje.

20.00 reception in the Jama hotel given by the President of
the Slovene Commission to UNESCO Prof. Dr. Peter
Tancig and the mayor of Postojna Mr. Josip Bajc

TUESDAY, June 28

- Lectures

8.00 Ognjen Bonacci, HYDROLOGY OF KARST SPRING
GRADOLE
8.30 Andrej Mihevc, RELIEF AT THE PONORS OF THE
KARST RIVERS - COMPARISON BETWEEN THE
CONTACT KARST AND KARST POLJES. THE
EXAMPLES FROM SLOVENIA
9.00 Jurič Kunaver, THE GEOMORPHOLOGY OF THE
TRANSITIONAL AND STEPPE-LIKE TYPE OF KARST
OF PODGORJE AND KRAŠKI ROB (KARST EDGE)
9.30 Paolo Forti, KARST HYDROTHERMAL SYSTEMS
10.00 Break
10.30 Branka Berce-Bratko, MAN AND KARST
11.00 Ines Ozboht, MENTAL PERCEPTION OF THE KARST
11.30 Bojan Žnidaršič, GEOGRAPHICAL NAMES IN
NOTRANJSKI PARK

13.00 Field work - Cerknjsko and Loško polje.

WEDNESDAY, June 29

Lectures

- 8.00 Andrej Kranjc, KARST POLJES MELIORATIONS AND REGULATIONS
- 8.30 Janja Kogovšek, SINKING RIVERS WATER QUALITY - PIVKA RIVER CASE STUDY
- 9.00 Tadej Slabe, SPELEOGENETICAL SIGNIFICANCE OF ROCKY RELIEF IN SOME CAVES OF NOTRANJSKA
- 9.30 Pavel Bosak, THE EVOLUTION OF KARST AND CAVES IN THE KONEPRUSY REGION (BOHEMIAN KARST, CZECH REPUBLIC)
- 10.00 Break
- 10.30 Andrej Gosar, APPLICATION OF GEOPHYSICAL METHODS IN KARST HYDROGEOLOGY AND ENGINEERING GEOLOGY
- 11.00 Daniel Rujšek, VELIKA VODA - REKA - A KARST RIVER
- 11.30 The conclusion of the Symposium
- 13.00 Field work - Notranjsko podolje and Rovte Contact Karst

THURSDAY, June 30

- 8.00 - 20.00 A whole-day excursion to karst poljes of Dolenjska

All the lectures will take place in Club Room of the Jama hotel

The departure to the field work and the final excursion will start from the Jama Parking place

On Tuesday and Wednesday evening there is the time at disposal for meetings of the international commissions and organizations or slides, videofilms etc. projection.

ABSTRACTS

MAN AND KARST

Branka Berce-Bratko*

The relationship between man and environment is basic for landscapes, either cultural or protected. The Karst as such is partially man's design created by use of the area.

Landscape definitions are evoking different images, and could be explained as visually-aesthetic element of the nature, as artist's image or an own pleasant experience, but less and less we associate the landscape with man's survival. Landscape is an expression of a place's regional context especially in the absence of distinguishing architectural styles. Regional context as its overall character, its geomorphology, water regime, soils and plants, animals and inhabitants represent regional "personality".

The research to this fundamental relationship emphasized identity. From anthropological identity to physical and social identity. A part of that was research on cultural landscape mapping, bioregionalism, and Risk assessment mapping as an example of MAB Reserve Buffer Zone Cerknjsko polje.

Villages around Cerknjsko plain and with this we became aware that all the research is only a memory, that being aware of past architectural values enables new guidelines for new developments. It represents an internal identity entered into the place, enhancing it and leaving signs of ancestors. High level of identity with landscape has enhanced the proper way of constructing, settlements design. This practice has enabled low level of erosion. The man and landscape are an entity. In the Notranjski Park area an old expression says: "everything is only returning, therefor such a research ought to be a base for new developments, like rehabilitation, re-instating genuine identity to people and places. This is not a conservation activity of the past but realization of past anticipations. The man is degrading places and mirrors its own limitations. The life is changing very quickly, settlements are loosing character and domestic appearance. Research of ethnoscares is research of people and landscapes, their inter-relationships and it will be shown on Cerknjsko polje, with particular reference to Žerovnica settlement and facing problems of eco-tourism. Special interest is education for environment and how to educate inhabitants and visitors to enhance protected landscape as a

* Urbanistični inštitut RS
Ljubljana (Slovenia)

part of MAB Programme NOTRANJSKI KRAS. Special presentation of the research of identity in its widest sense are geographical names in Notranjski park and Mental perception of the Karst as essential parts of the research study: MAN and KARST.

HYDROLOGY OF KARST SPRING GRADOLE

Ognjen Bonacci*

The paper discussed the results of a hydrogeological and hydrological analyses of the Gradole karst spring. Main goal of investigations was determination of catchment boundaries and area in order to protect spring water quality. The underground watershed has been determined by geological and hydrogeological methods. The control used was the hydrologic water budget analysis appropriate for karst basins with limited data. The catchment of Gradole spring is defined as 104 km². The Gradole spring water has been used as the drinking water supply. The capacity of the spring Gradole is limited and does not exceed 10 m³/s. All discharges which exceed this amount flow to the surface through other intermittent springs in the vicinity of Gradole spring. Measured minimum, average and maximum discharges in 1987-1992 period are: 0,28 m³/s, 1,80 m³/s, 8,68 m³/s respectively. During the hot and dry period of year there is a shortage of water for water supply. In this period an attempt to recharge the karst underground aquifer was made.

*Civil Engineering Faculty
Split University
Split (Croatia)

THE EVOLUTION OF KARST AND CAVES IN THE KONEPRUSY REGION (BOHEMIAN KARST, CZECH REPUBLIC)

Pavel Bosak*

Karst evolution. Lower Devonian - freshwater vadose and phreatic karstification connected with reef emergence and origin of freshwater lens - several phases in the interval of Pragian (Lower Devonian) to Givetian (Middle Devonian) - infill of clayey dolostone to dolomitic claystone, dark coloured - size millimetres to first tens of centimetres - depositional paleokarst. Pre-Cenomanian - not known subsurface karst forms - relies of surface forms and kaolinite sands in depressions. Paleogene - intensive karstification under the cover of Upper Cretaceous clastic sediments - intensive weathering of Cretaceous sediments - appearance of ponors - development of basic network of passages led to the origin of extensive caves - various fill, at the present time reworked completely. Miocene - gradual diminution of karstification - main periods of fossilization since upper Middle Miocene - speleothem origin - large lakes in caves - displacement by neotectonic movements. Pliocene to Early Pleistocene - continuing fossilization (Middle Pliocene and Biharian phases) - last neotectonic movements during Biharian (cca 11 Ma). Pleistocene to Holocene - intensive backward erosion at the end of Biharian - superficial karstification (epikarst zone) - maybe inserted slope caves - karstification up to local base level.

Paleohydrogeological Model. (1) Predisposed network of Lower - Middle Devonian diagenetic porosity - several times utilized by all younger karstification phases. (2) Evolution of some karstic porosity to macroporosity during pre-Cenomanian evolution of landscape. (3) Destruction of platform cover of Upper Cretaceous sediments under wet and hot climate (Paleocene, Eocene, Oligocene) and (*) evolution of ponors in multiple ranks and multiple lines at margins of limestone core of synclines. (*) evolution of infiltration routes under a cover of permeable Cenomanian sediments. (4) Evolution of cave systems - deep phreatic cave horizons with multiple loops and local horizontal features - followed main tectonic scheme - NNE-SSW to NNW-SSE tension open lines and WNW-ESE to WSW-ENE compressional faults to overthrusts - main characteristics: - in areas close to ponors - circular,

*Czech Speleological Society
Praha (Czech Republic)

mostly phreatic tunnels, increasing diameter with increasing distance from ponors, - in area of main hydraulic barrier - irregular maze to labyrinth caves, vertical with subhorizontal "levels" - morphology of shallow phreatic to vadose lake-filled rooms. (5) Main hydraulic barrier - the Očkov Overthrust - upwell of water - increase in piezometric level - water discharged by this thrust line in NW and SE direction. (6) Evolution of lower deep phreatic cave horizons in dependance to decrease of piezometric level - ?owing to climate aridization and/or ?deepening of regional base level - maybe uplift connected with volcanic phases - lowering of piezometric level marked by precipitates of Fe inside the carbonate massif - retreat of limestone slopes and shift of ponors in the centripetal manner.

STRUCTURE OF IDRIJA FAULT ZONE ON PLANINSKO POLJE Jože Čar*

The Idrija fault is one of the strongest and morphologically the most distinctive tectonic zone in the area of Southern Alps. It passes from the Rečija valley, cuts the whole western Slovenia and continues at Prezid in Croatia. In extremely complicated wider fault zone of the Idrija fault there are big karst poljes, namely Planinsko and Cerkljsko polje, Loška dolina and Bahno polje. The wider fault zone was mapped in detail on Planinsko polje and its vicinity in particular. The Idrija fault is a wrench of continental character this is why in its inner and outer and wider fault zone richly developed various forms of broken, collapsed and fissured zones having strong influence on polje's origin and on its hydrological properties.

*FNT, VTOZD Montanistika,
odsek za geologijo
Ljubljana

KARST HYDROTHERMAL SYSTEMS

Paolo Forti*

Thermal karst systems may represent an important economic resource from the point of view of the hydrogeology and of the hosted ore deposits. Nevertheless general papers on this topic are very scarce in the international literature and completely lacking in the Italian one.

In the past every time a cave showed strange morphologies or was infilled by exotic deposits like uncommon speleothems or several different minerals it was normal consider it developed under hydrothermal conditions.

In reality, in the last years the improvement of our knowledge on the different and time to time more complex speleogenetic mechanisms demonstrated that several previously supposed hydrothermal karst system should be developed in a normal environment.

In theory a cave may be regarded to as hydrothermal in origin (or part origin) if thermal fluids (normally hot water) flowed inside it (or part of it) at least during a period of its development.

Even if this definition is very simple the criteria by which a cave may be regarded as thermal are rather complicated and often non conclusive: the most certain of which (though uncommon) is hot water flowing inside the cave.

In the literature several papers exist on peculiar thermal karst systems, but very few of them give a general overview on this topic (Ford & Williams, 1989).

Moreover most of the research deals with the morphology of these cavities. From this point of view the hydrothermal caves may be subdivided into two categories (Muller & Salvary, 1977): monogenic hydrothermal caves, consisting of a large basal chamber from which a branching pattern of rising bell shaped passages have grown; the second category consist of 2D- and 3D- maze caves. The possibility of evolution of the different types of hydrothermal caves depend upon the fracturation of the rocks, the presence of an overlying watershed, the kind of feeding and discharge and other parameters (Muller & Salvary, 1977; Dublinsky, 1980; Bakalowiez et Al., 1987).

*Istituto Italiano di Speleologia
Bologna (Italy)

Until now only a few papers have been produced on the peculiar speleothems growing in the hydrothermal caves and none of them has a general approach (Hill & Forti, 1986); on the contrary a wide bibliography exists on the ore deposits (Mississippi Valley Type) related with thermal karst systems (Dzulynsky & Sass-Gustkiewicz, 1989).

A general study was never performed to relate morphologies and chemical deposits of the thermal karst with the speleogenetic mechanisms which may be active in the different zones of the aquifer.

In reality due to the complexity and the variability of the thermal fluids, there are plenty of different chemical-physical reactions which are possibly active in such an environment; anyway some of them may be regarded as the most important, because they are active singularly or combined with some others at least in part of most of the hydrothermal caves.

Some of them cause net dissolution, or net deposition, but most of them are responsible for incongruent dissolution (deposition and dissolution in the mean time). Normally these mechanisms may be active only in a definite part of the aquifer and therefore their effects are possibly indicators of the conditions in which the cave developed.

Therefore it would be extremely important to define the interrelations existing between speleogenetic mechanisms, aquifer zones and evolved morphologies and/or chemical deposits: in the present paper a first schematic attempt of such classification is proposed for discussion.

References

- Bakalowicz M. J. & Ford D. C. & Miller T. E. & Palmer A. N. & Palmer M. V., 1987: Thermal genesis of solution caves in the Black Hills, South Dakota, *Bull. Soc. Geol. Am.*, 99, 729-738
- Dubiansky V. N., 1980: Hydrothermal karst in the alpine folded belt of southern part of the U. S. S. R., *Kras i Spel.* 3 (13), p. 18-36
- Dzulynsky S. & Sass-Gustkiewicz M., 1989: Pb-Zn ores in Bosak P. ed. "Paleokarst", Academia, Praga, p. 377-393
- Ford D. & Williams P., 1989: Karst geomorphology and hydrology, Unwin Hyman, London, p. 282-293
- Hill C. A. & Forti P., 1986: Cave Minerals of the World, NSS, p. 1-238
- Muller P. & Salvary I., 1977: Some aspect of the developments in Hungarian speleology theories during the last 10 years. *Karst-es-Barlang*, p. 58-59

TYPES OF POLJES IN SLOVENIA AND THEIR INFLUENCE ON LAND USE

Ivan Gams^{*}

The poljes bottom in Slovenia are at the most used on the gravel, sand and thick loam soil on the alluvial plains and the least on the shallow stony rendzina and flooded areas. The deposition of river load is mostly connected with the polje types (border, piedmont, peripheral, overflow, base level polje). For instance, in the border and piedmont polje bottom is more frequent the gravel and sand deposition and floods are often in the base level and overflow polje. These correlations are shown in the case of the poljes in the Dinaric Karst and Alps in Slovenia. The largest poljes belong to the combined types.

APPLICATION OF GEOPHYSICAL METHODS IN KARST HYDROGEOLOGY AND ENGINEERING GEOLOGY

Andrej Gosar^{**}

In karst areas, regional geophysical surveys are centred on the assessment of karstified permeable zones, verification of the structural-tectonic position of limestones with respect to their basement and determination of the spatial distribution of carbonate rock complexes. Regional gravity and (aero)magnetic maps are useful for studying the relief of impermeable basement rocks, whose dips control the general directions of ground-water flow in a limestone block and for location of tectonic lines. Zones of intensive karstification are usually associated with faults and especially with intersections of tectonic systems.

Detailed geophysical methods are applied to study surface (sinkholes, depressions) and underground (caverns, ground-water conduits, underground streams) karst formations. Because these forms appear as electrical more conductive zones concerning adjacent undisturbed

^{*}Pohorskega bataljona 185

Ljubljana

^{**}Institute of Geology,

Geotechnics and Geophysics,

Ljubljana (Slovenia)

limestones, geoelectrical methods are the most important. Vertical electrical sounding (VES) provides information on intensity of karstification in a vertical section that is normally composed of four principal layers: near surface clayey sediments, dry karstified limestone, water-bearing karstified limestone and solid unkarstified limestone. Direction of ground-water flow can be determined with *mise-à-la-masse* method, measurements of spontaneous potential (SP), very low frequency (VLF) electromagnetic method and with geothermal methods. Induced polarization (IP), VES and SP methods provide valuable information about the overburden sediments that control the infiltration of surface water into limestones and about bauxite deposits.

A special task for geophysics on karst is locating and tracing cavities (channels and caves). If they are filled with water, geoelectrical methods are applied as in tracing conductive zones. In case of dry, isolated cavities microgravimetry or ground penetrating radar (GPR) is used. Fractures and karstified zones in general are investigated also by reflection and refraction seismics and gravimetry combined with resistivity methods.

In Slovenian karst researches from Institute of Geology, Geotechnics and Geophysics from Ljubljana successfully applied geophysical methods for several times in solving different problems including: - determination of the base of karstification with vertical

electrical sounding, - tracing of ground-water flow with SP, *mise-à-la-masse* and

geothermal methods, - surveying of bedrock (sinkholes, depressions) on karst fields

with resistivity profiling, - investigations of karstified tectonic zones (also under

flysch deposits) with reflection and refraction seismics

combined with vertical electrical sounding, - location of caves with ground penetrating radar and

gravimetry, - studying of bauxite deposits with different geoelectrical methods.

SINKING RIVERS WATER QUALITY - PIVKA RIVER CASE STUDY Kogovšek Janja*

The paper deals with the water quality of the Pivka sinking stream at its ponor to Postojnska jama. On its superficial way towards the ponor Pivka receives the waste communal and industrial waters of the villages, only the waste waters of Postojna are being treated. By various methods of sampling we tried to encompass the majority of properties of Pivka, its water quality respectively. Seasonal observations evidence the oscillations of particular parameters during the observations and specially called the attention to critical situations which occur during winter and summer drought. They are conditioned by low water level and sinking of Pivka in upper part of its bed. The hazard is reflected in the content of dissolved oxygen, COD and BOD. Detailed systematic observations during 24 hours gave the quality properties during low winter and summer water level as well at increasing and decreasing of the discharge.

The Pivka quality downstream, in Pivka and Planinska jama depends on the water quality at the ponor. Known and unknown underground tributaries with pollution or dilution effects, autopurification processes active on the Pivka underground flow have impact on this quality.

KARST POLJES MELIORATIONS AND REGULATIONS Andrej Krauž*

With the economical and technical development pressure of agrarian population on farming land, including the wishes to protect the cultivated land against floods and to cultivate new lands, increased at the end of 18th Century. Such trends were specially strong on the karst poljes with larger flat soil bottom and at the same time more often flooded. During 19th century more plans to prevent floods on Cerkljsko polje primarily were elaborated.

* Institut za raziskovanje kraša ZRC SAZU
Postojna

In 1886 the Ministry for Agriculture entrusted to V. Putick the preparation of the projects to abolish the floods on the karst poljes of Carniola. Putick made a project "Generalproject zur unschädlichen Ableitung der Hochwässer aus den Kesselthälern von Planina, Zirknitz und Laas-Altenmarkt in Innerkrain" (1888), but it was not adopted. Nevertheless some proposals were realized under his direction already, and some of them much later. Most of the visible meliorations on our karst poljes are in this way or that way connected with Putick's activity. Putick's influence was large enough and he was asked to make plans and even to realize them on some other karst poljes in Slovenia too. In the meantime other engineers were recruited to make the meliorations on other poljes, specially in the background of Krka river springs. During the 20th Century plans for the meliorations of karst poljes included also the plans to make water accumulations for HE power stations (Planinsko polje) or to change them into permanent lakes (Cerkniško polje).

THE GEOMORPHOLOGY OF THE TRANSITIONAL AND STEPPE-LIKE TYPE OF KARST OF PODGORJE AND KRAŠKI ROB (KARST EDGE)

Jurij Kunaver¹

The region of the southwestern Slovene Dinaric karst plateaus which is represented with Podgorski kras as the biggest area has a typical transitional and steppe-like nature, from higher to lower karstic plateaus and finally to the flysch region, along the coast in the western Slovenian Istria. The main character of the country is the frequent interruption of the low lying karstified plateaus with vertical structural steps or escarpments. The area is therefor also called Kraški rob (The Karst Edge). It extends from Socerb on NW to Golič on SE and in the altitude from about 400 m to nearly 900 m, with the highest point on Slavnik, 1001 m. This block which belongs to the Čičarija imbricate structure, has a thrusting geological structure and intercalations of the flysch between the limestone thrust folds and nappes. Typical are the elongated limestone stripes which are interrupted by very narrow stripes of flysch.

¹Oddelek za geografijo FF
Aškerčeva 2
Ljubljana

Geologically the region is the youngest part of the westernmost Dinaric geologic belt with the transitions from Lower to Upper Cretaceous and in continuation from Lower to Middle Paleocene to the Upper one and to Eocene flysch. The surface of the karst plateaus shows a clear evidence of the recent, presumably Pleistocene and partly Holocene river drainage and has a fluvial karstic nature. There is a clear evidence of a continuously changing relation between the karstic and flysch regions, the later being under the influence of more dynamic erosional and denudational processes and therefore being a subject of the territorial diminishing. The karst area as a whole has a shallow character lacking deeper karst depressions. The shallow dolines mostly transformed by man express frequency of up to 54/km². Some parts of the surface are stony but the reforestation and abandoning of the intensive agriculture cause the changing of the landscape. Except of the plateaus where the flysch bases are used for settlements and fields, the limestone walls between them with concave rock shelters or abris on their bottom exposed to SW, are the second most impressive feature in the region. The walls which are the result of recumbent folds, continue from the Triest Karst along the border with flysch to Croatian Istria. Except the rock shelters also the caves are to be found near the thrust plane on the contact of the limestone rock and the flysch at the base of the walls.

RELIEF AT THE PONORS OF THE KARST RIVERS -
COMPARISON BETWEEN THE CONTACT KARST AND KARST
POLJES. THE EXAMPLES FROM SLOVENIA
Andrej Mihevc^{*}

Along allogenic rivers, flowing to the karst, bigger depression forms developed, usually called blind valleys. More accurate view shows that they differ among them considerably. Various depression forms may be morphologically distinguished and genetically founded. These are the ponor steepheads, ponor valleys, blind valleys with corrosionally widened bottom. These shapes originated under the influence of the fluvial inflow but they are controlled by hydrological properties of the karst where they sink.

^{*} Inštitut za raziskovanje krasa ZRC SAZU
Postojna

Karst poljes are bigger depression forms where superficial streams exist. The corrosionally levelled karst poljes bottoms are formed at the level of karst water. At the ponors the impact of bigger karst water gradient is shown evidencing the property of the karst lying below the ponors.

This is why at the ponors on karst poljes the relief forms develop that are similar to the forms of the contact karst.

MENTAL PERCEPTION OF THE KARST

Ines Ozholt*

From the work in the Project of Notranjski Park the difference between general perception in a certain area and the perception of the same area on individual level is clearly evident.

Where spatial management or planning is concerned the mental image is merely ignored, although it is well known that this particular image is of vital importance for the people living in the area. This is the main reason that we propose the preliminary study of perception of the place from the point of view of inhabitants and visitors before any of the planning procedures take place.

The perception is one of the basic functions of man's orientation in a place and has great impact. Its importance orientation in a place and has great impact. Its importance should be considered in the procedure of physical planning. The identity of territory has impacts on visual and sensual perception of a specific area. The area which is heterogeneous has its identity which has a higher degree of recognition for people than an area lacking variety.

The Notranjski Park has great diversity of landscapes and is interesting either for research or exploration in a view of professional responsible for the establishment of the park. The project begins with the testing of the Karst landscape. One method is used for the entire area of the Notranjski park. We intend to determine, whether the Karst, as a land, has an identity of it's own, which is different from the identity of other regions both in the natural and social point of view. We intend to determine, whether the people of this region differ their home from other landscapes both visually and mentally. It also concerns us, how Karst is interpreted by the natives and the tourists, and what are the differences between the impressions of professionals and "laics".

*Arca Pro d.o.o.
Cerknica (Slovenia)

The participants of the International Karstological School will be asked to fill in questionnaires on recognition and mental image of the Karst. They will represent the first pilot group, the one representing professionals, second group are visitors, the third are indigenous people.

Tasks of the project:

- to establish the existence or non-existence of difference between mental image of the Karst phenomena and other geological features and areas,
- to establish which are basic expressions interconnecting the Karst as such and the Karst phenomena,
- to establish the mental image of the Notranjski Park according to their professional experience and the knowledge,
- to evaluate the Karst as the place of pleasant or non-pleasant experience,
- to establish which elements of the Karst are the most important for its recognition; elements of macro-reliefs or micro-reliefs,
- to establish the existence or non existence of differences between mental images among three pilot groups.

Methods used:

1. "knowledge questionnaire" (open and closed questions).
2. scales of measures for cognition (semantic differential).
3. association tests.
4. photo-questionnaire.

Sample:

Group of professionals (people researching or studying Karst).

Group of visitors (enthusiasts, tourists enjoying the Karst).

Group of indigenous people (people living in the Karst area perceiving it as home environment).

Procedure:

The results will be reached by questionnaire, which is the same for all pilot groups with contents: knowledge, association, perception, recognition, and assessments.

Results:

Mathematical and statistical methods will be used to generate research result and to establish the level of achieved aims (factor analysts, variance analysis with F test used).

VELIKA VODA - REKA - A KARST RIVER

Daniel Rojšek**

Velika voda - Reka river is known in natural sciences and literature from antiquity. In the Postojna Karst Institute rich bibliography of 189 units exists.

Hydrogeographical features of the Velika voda - Reka fluvio-karst basin have been studied and published, but the river regime of 30 years period (1953-1982) has been proceeded for this occasion.

Hydrogeographical characteristics of the Velika voda - Reka fluvio-karst drainage area and of the matični Kras flowing-through karst area will be presented in the communication. Hydrogeographical heritage of the area will be mentioned, too.

SPELEOGENETICAL SIGNIFICANCE OF ROCKY RELIEF IN SOME CAVES OF NOTRANJSKA

Tadej Slabe***

Studying the origin and formation of the cave rocky forms it was proved that combined into rocky relief they are frequently an important speleogenetical factor. This cognition is confirmed by study of rocky relief in selected caves of Notranjska. The results may be used to define the origin and development of this morphogenetically joint karst area (Fig. 1). I have chosen the caves at the inflow (Križna jama) and outflow (ponor caves Mala Karlovica and Zelške jame) sides of the Cerkljeva lake and active effluent cave Planinska jama where the waters of Pivka and Rak flow together. At NW, higher part of Notranjska lowered surface, it means at the border of Črni vrh polje (Gams 1974) I have studied Ciganska jama near Predgrize. Logaska jama is an old cave lying on Logatec karst plain.

*Zavod za varstvo naravne in kulturne dediščine
Nova Gorica (Slovenia)

***Inštitut za raziskovanje krasa ZRC SAZU
Postojna

POLJES AND CAVES OF NOTRANJSKA France Šusteršič*

The Eastern (Cerknica) branch of Ljubljana basin encompasses four well pronounced poljes, and additional five closed basins which display at least some properties of poljes. Though similar by their origin the poljes display different morphology which is due to local conditions. The same goes for the caves. Among them appear a number of caves which cannot be related to the present hydrological situation.

GEOGRAPHICAL NAMES IN NOTRANJSKI PARK Bojan Žnidaršič**

As the basis for the research in the area of Notranjski Regional Park computer aided data base was formed in which available geographical names were used a base. All parameters important for the context and statistics of geographical names were analytically assessed.

Particular names ascribed by people were either for the purpose of orientation or rational-logical organisation of space. For the purpose of this particular research project two divisions were made:

- visual (structure and relief)
- substance (structure and quality).

The most important characteristics (i.e. use of place) are evident in place names. Nomination of names for certain parts have different background as they have continuing history and could be ascribed to different circumstances as well. Geographical names represent the use of the place, predominantly its use in the past. As such they are a part of a cultural heritage.

In the introduction the purpose of the research was presented as well as the hypothesis. The applicability of such kind of the research in different levels was evaluated. In the disposition of the research the

*FNT, VTOZD Montanistika,
odsek za geologijo
Ljubljana
**Area Pro d.o.o.
Cerknica (Slovenia)

contents of inventory phase and analytical levels chosen were justified. Following the description of the procedure the detailed structuring of parameters was presented. The method of data collection and assessments pursued were described precisely. The instructions are given for relations and connections of geographical names in the data base.

All important parameters are evaluated (there are 72 parameters). 11 content, structures:

- level of altitude,
- exposition,
- use,
- partitions,
- peaks,
- valleys,
- waters,
- microrelief features,
- routes,
- settlements,
- objects - buildings.

The main part of the research consists of contents analysis by statistical assessments. The findings are shown in graphs and charts, in five charts all available geographical names (5392) are stated, different names of places (4446) and the difference between those two categories (946). The Chart of most common place names and the chart of "naming" diversity of places was produced (149). With the help of 14 graphs the percentage of each parameter were presented.

The summary of inventory work used as the basis for the continuation of the research - second phase definition. Three new substances were suggested: spatial, socio-economic and linguistic. In the first phase one of the possible structural results is presented.

For the one of the possible structure research the study of the impact for the karst phenomena on naming geographical places. Therefore for the second International Karstological School the research results are summarized, and geographical names stemming from karst phenomena are presented. As a starting point was a question: where, how much, and with which elements of the area studied the karst phenomena appear as the basis for naming the places. This way 306 names based on karst phenomena were identified.

The study has appendix supporting the transparency of the approach, summary and literature used, where all available bibliography is collected. The aim of the study was to put together all available knowledge, experiences, and achievements of all authors dealing with this particular problem of geographical names in Slovenia.

Field-work Schedule

No special equipment is required for the field-work. We shall travel by bus and walk only 2 km at once. The only inconvenience that threatens may be the summer storms.

MONDAY, June 27: **PLANINSKO POLJE**

Departure at 13.00

Entrance to Planinska jama, steephead valley, spring of Majni and steephead valley, Laze village and ponor side of polje, ponors at Milavčevi ključi, Idrija fault line and connected tectonics, ponors Pod Stenami.

19.30 return to Postojna

TUESDAY, June 28: **CERKNIŠKO POLJE**

Departure at 13.00

Ponor side of polje, entrance of Mala and Velika Karlovica caves, ponors at Vodonos, springs Obrh and Cemni, Loško polje, entrance to Golobina cave, spring of Žerovnica, surface river Cerkniščica at Bloke.

20.30 return to Postojna.

WEDNESDAY, June 29: **HOTEDRŠICA AND LOGATEC BORDER POLJES**

Departure at 13.00

Border of Planinsko polje, Hotenjski ravnik flat corrosional surface, Hotedršica alluvial fan, Logatec border polje, Route contact karst, springs of Ljubljaničica at Vrhnika and Bistra.

20.30 return to Postojna.

THURSDAY, June 30: **KARST POLJES OF DOLENJSKA**

Start at 8.00, return at 20.00

Ljubljana Moor, Grosuplje basin, Radensko polje, Čusperk, Ilova gora, Ciganska jama, Zagradec, Mlačevo, Žalna, Žalnsko polje, Luče, Škrjanceva jama (catastrofhe), val clos and Krke resurgence, Dobrepolje, Ribniško polje, Tentera (cave), Ribnica, ponor of Bistrica river, Rakitnice resurgences, Rakitnica ponors.

Lunch time

Kočevo polje, caves at Klinja vas, Željinc, Ciganska jama, Kočevje, Livold, Kočevska reka, Glažuta, Loski Potok, Blaško polje, Cerknica, Postojna.

20.00 return to Postojna.

The sources from which the annexes to the field-work are used:

- Čar, Jože & Gospodarič, Rado, 1984: About Geology of Karst among Postojna, Planina and Cerknica.- *Acta carsologica*, 12 (1983), 93-106, Ljubljana
- Gams, Ivan, 1981: Inundations in the polje of Planina.- *Geografski zbornik*, 20, 9-29, Ljubljana
- Gospodarič, Rado & Habič, Peter /Editors, 1976: Underground water tracing, investigations in Slovenia.- Karst Research Institute, 1-312, Ljubljana
- Gospodarič, Rado & Habič, Peter, 1970: Karst phenomena of Cerkniško polje.- *Acta Carsologica*, 8, 1, 11-162, Ljubljana
- Gospodarič, Rado, 1970: Speleological investigations of the Cerknica cave system.- *Acta Carsologica*, 8, 111-169, Ljubljana
- Gospodarič, Rado, 1973: Viršnica - Der Höhlensystem von Šica am Radensko polje.- *Naše jame*, 14(1972), 25-33, Ljubljana
- Guide-book of the Congress Excursion through Dinaric Karst, 1965.- 4. Congrès International de Spéléologie, 1-201, Ljubljana
- Habič, Peter & Kogovšek, Janja & Bricej, Mihael & Zupan, Martina, 1990: Dobličica springs and their wider karst background.- *Acta carsologica*, 19, 7-100, Ljubljana
- Habič, Peter, 1985: Water table in Slovene karst of Notranjsko and Primorsko.- *Acta carsologica*, 13, 37-78, Ljubljana
- Habič, Peter, 1987: The ponor cave Golobina on Loško polje.- *Acta carsologica*, 16, 37-49, Ljubljana
- Hrovat, Alojzij, 1959: Le poljé près de Žalna.- *Naše jame*, 1, 2, 53-58, Ljubljana
- Kranjc, Andrej A., 1982: Historical methods of water investigations in Dolenjsko (Slovenia).- *Naš krš*, 6, 12-13, 3-10, Sarajevo
- Kranjc, Andrej, 1981: The karst development in "Ribniška Mala gora" (Slovenia, Yugoslavia).- *Acta Carsologica*, 9, 31-85, Ljubljana
- Melik, Anton, 1955: Les poljés karstiques de Slovénie au pléistocène.- *Razred za prirodoslovne vede, SAZU*, 7, 1-162, Ljubljana
- Melik, Anton, 1959: Posavska Slovenija.- *Slovenija, Geografski opis, Slovenska Matica*, II, Tretji zvezek, 9-695, Ljubljana
- Meze, Drago, 1981: Flood areas in the Grosuplje basin.- *Geografski zbornik*, 20, 37-92, Ljubljana
- Novak, Dušan, 1974: Sur la hydrographie du poljé de Kočevje.- *Acta Carsologica*, 6, 369-394, Ljubljana
- Novak, Dušan, 1987: Podzemeljski vodni tokovi na Dolenjskem.- *Dolenjski kras*, 2, 23-27, Novo mesto


Routes of the excursions-


1. Planinsko polje

2. Cerkljско polje


3. Hotedratska and Logatec

4. Karst poljes of Dolenjska





Karta 1



Sl. 14. Razporeditev nizkih voda v Notranjskem krasu
 1 — kraška polja in dolina Ljubljanskega barja, al
 2 — fliš, nepropustno površje
 3 — kras, apnenec in dolomit
 4 — hidrokorrelacijske nizke vode v krasu
 5 — površinsko razvodje
 6 — kraško razvodje
 7 — kraški izvir in ponikovalnica

Fig. 14. Distribution of low waters on Notranjsko karst
 1 — karst poljes and basin of Ljubljana Moor, allu
 2 — flysch, impermeable surface
 3 — karst, limestone and dolomite
 4 — hydro-correlation-lines of low water in karst
 5 — superficial watershed
 6 — karst watershed
 7 — karst spring and sinking stream




Fig. 5. Schematic review of superficial and underground waters in Ljubljana karst river basin. 1 — important gauging stations, 2 — karst polje with sediment, 3 — central hydrogeologic relative barrier, 4 — intake area of permanent karst springs, 5 — periodic high water discharges of karst water, 6 — underground water connections, 7 — superficial stream with springs and ponors, 8 — superficial watershed, 9 — supposed karst watershed.

SL 5. Shema površinskih in podzemeljskih voda v kraškem porečju Ljubljane. 1 — važnejše vodomerne postaje, 2 — kraško polje z naplavinami, 3 — osrednja hidrogeološka pregrada, relativna bariera, 4 — zbirno območje stalnih kraških izvirov, 5 — občasni visokovodni prelivi kraške vode, 6 — podzemeljske vodne zveze, 7 — površinski tok z izviri in ponori, 8 — površinsko razvodje, 9 — kraško predpostavljeno razvodje.

Fig. 2. Longitudinal cross section of central part of Ljubljana karst river basin. 1 — active cave or polje, 2 — unknown parts within known active caves, 3 — the supposed zone of former active caves between Planinsko polje and Ljubljansko barje (Ljubljana Moor), 4 — karst polje's bottom, 5 — Gradšče relief level, 6 — Bodiški vrh relief level, 7 — Dolina relief level.

SI. 2. Vzdoljni presek osrednjega dela kraškega porečja Ljubljanske. 1 — vodna jama ali brezno, 2 — neznan odsek v znanih vodnih jamah, 3 — predvidena zona nekdanjih vodnih jam med Planinskim barjem in Ljubljanskim barjem, 4 — dno kraškega polja, 5 — Gradšče reliefna stopnja, 6 — Bodiška reliefna stopnja, 7 — Dolinska reliefna stopnja.




Fig. 6






Fig. 8. Low and high water gradients between Pivka basin and Planina polje


Fig. 7



POSKA KARTA
OLOGICAL MAP



2 km







Fig. 9






Fig. 18. The situation and cross section of caves at eastern Planinsko polje border. 1 — karst polje's border with superficial river bed and swallow-holes, 2 — collapsed dolines, 3 — simplified caves ground plan, 4 — lowest known karst water level — hydraulic contour lines, 5 — direction of karst water outflow, 6 — the floods altitude on polje and in caves, 7 — the pools of caught water in the cave, simplified cave's cross section.

Sl. 19. Položaj in prerez jame ob vzlednem robu Planinskega polja. 1 — rob kraškega polja s površinsko strugo in požiralniki, 2 — udornice, 3 — poenostavljeni tloris jame, 4 — najnižja znana gladina kraške vode — hidroizokliza, 5 — smer odтока kraške vode, 6 — višina poplave na polju in v jamah, 7 — bazeni ujete vode v jami, poenostavljeni prerez jame.

Fig. 11






Diagram 1. Trajanje pretokov in mesečni pretoki Unice - Hasberk v 40-letnem razdobju
 Diagram 1. Duration of Discharge and Monthly Discharge at Hasberk in 40-years Period.

Trajanje pretokov Unice - v. p. Hasberk



Mesečni pretok Unice - v. p. Hasberk



Vr. Poročila SUOVI 1974

Geografska inštitucija Anzonia Melka SAZU *

Tabela 1. Povprečna letna števila dni z vodno gladino v. p. Unica-Hasberk (1949-1977)

Table 1. Average yearly number of days with water level at the gauge station at the Unica-Hasberk (1949-1977)

Mesec	Višina vodne gladine – Altitude of water level						Skupno > 447,0 m
	< 447,0	447,0– 448,0 m	448,0– 449,0 m	449,0– 450,0 m	450,0– 451,0 m	451,0– 452,0 m	
I	26,6	3,2	1,2				4,4
II	25,1	2,8	0,3				3,1
III	27,0	1,7	0,3				4,0
IV	24,7	3,5	0,7	0,5	0,5	0,1	5,3
V	28,8	1,7	0,1	0,2	0,2		2,2
VI	29,2	0,8					0,8
VII	30,8	0,2					0,2
VIII	30,6	0,4					0,4
IX	27,6	2,0	0,4				2,4
X	26,8	3,6	0,4	0,2			4,2
XI	73,6	5,1	1,2	0,1			6,4
XII	23,1	6,1	1,7	0,1			7,9
Letno	373,8	43,1	6,3	1,1	0,7	0,1	41,3

Tabela 2. Padavine i potencijalna evapotranspiracija u Planini (1931-1960).

Table 2. Precipitation and potential evapotranspiration at Planina (1931-1960).

Mesec Month	Padavine, mm Precipitation mm	Potencijalna evapotranspiracija Potential evapotranspiration mm	Razlika Difference
I	131	0	131,0
II	126	0,8	125,2
III	119	17,3	101,7
IV	126	40,7	85,3
V	130	78,7	51,3
VI	163	104,8	58,2
VII	150	134,8	15,2
VIII	137	103,7	33,3
IX	168	72,8	95,2
X	194	42,3	151,7
XI	184	21,2	162,8
XII	179	3,0	167,0
I-XII	1831	600,2	1224,9
IV-IX	894	515,5	378,5

Tabela 3. Mesečni koeficijenti odтока padavinske vode, protoka Unice u njenom vodostaju nad 446 m
Table 3. Monthly coefficients of drainage, flow of the Unica and its water level above 447 m (the highest value = 12)

Mesec Month	Koeficijent odтока padavinske vode (1930-1960) Coefficient of the drainage (1930-1960)	Koeficijent mesecnog protoka Unice (40 let) Coefficient of the flow of the Unica (40 years)	Vodostaj Unice nad 447 m Water level of the Unica 447 m (1944-1977)
II	8	5	6
III	7	9-10	7
IV	5	9-10	10
V	4	6	4
VI	3	4	3
VII	2	2	1
VIII	1	1	2
IX	6	3	5
X	10	7	8
XI	11	12	11
XII	12	12	12

Tabela 4. Vodostaj Unice pod mostom pri Hrabniku

Table 4. Water level of the Unica below the bridge at Hrabnik

Vodostaj - Water level	Razdoblje - Period	Letno brojilo dni Yearly number of days
< 447 m	1970-1977	107,4
	1949-1977	123,9
447-448 m	1970-1977	38,7
	1949-1977	33,0
448-449 m	1970-1977	12,0
	1949-1977	6,6
449-450 m	1970-1977	2,7
	1949-1977	1,2
450-451 m	1970-1977	2,5
	1949-1977	0,7
451-452 m	1970-1977	0,2
	1949-1977	0,1






Fig. 15



KARLOVICE




Planis izid ob potočni strani Ceratnikskega polja


1 — skala na vodu 2 — kamen in šot 3 — podoben skale 4 — tla iz glin 5 — kapnik 6 — rešer ob nizki vodi 7 — vodni strugi in brzice 8 — sifon 9 — smer vodnega toka 10 — vodni strugi na površju 11 — skalni rob polja 12 — smer in vpad skladov 13 — prelomi 14 — poligonske točke 15 — A-A' in B-B' so prečni prerezi na št. 2

The Ground Sketch of the Caves at the Caves at the Poma Side of the Ceratnik Polje

1 — the rocky floor in the channels 2 — logs and stree 3 — breakdown rocks 4 — the floor covered with water 5 — formations 6 — lakes at low water 7 — the water bed and rapids 8 — ghyben 9 — the direction of water current 10 — the river bed on the surface of the polje 11 — the rocky edge of the polje 12 strike and dip of beds 13 — fault 14 — polygonic ponds 15 — A-A' and B-B' are cross sections in fig. 2



Sl. 7. Smeri kraške vode med Cerklješkim jezerom in Planinskim poljem
Fig. 7. Karst water gradients between Cerklje lake and Planina polje




Sl. 10. Naplavinne med Dolnja jezero in strugo Stržena upoštevane z podrobno vrstnami V1 — V12; podrobnejši opis plastí glej v tekstu

1 — humus, 2 — rjavkasta ilovica s hitinami polžev, 3 — siva ilovica s hitinami polžev, 4 — dolomitni grušč, pesek in siva ilovica, 5 in 6 — rumenkasta ilovica in droben kremenov pesek, 7 in 8 — rjava peščena ilovica, pesek, prod in prepečen grušč, 9 — pasovita sivorjava ilovica s peskom, A — mlajši zasip, B — jezerski sedimenti, C — starejši zasip

Fig. 10. Sediments between Dolnja jezero village and Stržen creek bed, based on bore-holes V1 — V12; detailed description of the layers in the text

1 — humus, 2 — brownish loam with snail shells, 3 — grey loam with snail shells, 4 — dolomite rubbles, sand and grey loam, 5 and 6 — yellowish loam and silt, 7 and 8 — brown sandy loam, sand and gravel, weathered rubbles, 9 — greybrown varved loam with sand, A — younger fluvial fill, B — lake sediments, C — older fluvial fill




Sl. 28. Morfologija Vodnosa.

1 — stari ugrezi v okolici ponikve, 2 — novi ugrezi v okolici ponikve, 3 — grezi v ponikvi, 4 — razkrita skalna podlaga, 5 — obod ponikve

Fig. 28. Morphology of Vodnos ponor

1 — old sinkholes in the ponor vicinity, 2 — recent sinkholes in the ponor vicinity, 3 — sinkholes in the ponor, 4 — exposed rocky bottom, 5 — ponor boundary




Sl. 1. Položaj Golobine in požiralnikov na Loškem polju

a — jama, b — umetni kanal, c — občasni požiralniki, d — stalni požiralniki, e — skalni obod polja

Fig. 1. Golobina and swallow holes situation on Loško polje

a — cave, b — artificial channel, c — periodical swallow holes, d — permanent swallow holes, e — rocky polje's border

Priloge 1. Posredna zona Golobina pri Danah, Uroš
 Annex 1. Posredna zona Golobina pri Danah, ground plan







Fig. 2. Sinking streams and dry valleys on Logatec-Rovte plateau. 1 — superficial watershed, 2 — sinking stream with the ponor's sea level, 3 — dry valley, 4 — Quaternary sediments on marginal karst polje, 5 — Dolina or Ravnik relief level, 6 — Rodiški vrh level, 7 — Gradšče level.

Sl. 3. Ponikalnice in suhe doline na Logaško-rovteški planoti. 1 — površinsko razvodje, 2 — ponikalnica z nadmorsko višino ponora, 3 — suha dolina, 4 — kvartarne naplavinne na robnem karškem polju, 5 — Dolinska ali Ravniška reliezna stopnja, 6 — Rodiška stopnja, 7 — Gradšča stopnja.

Fig. 24



Sl. 4. Stopni nizkih in visokih voda med Planinskim poljem in izviru Ljubljance.
Fig. 4. Gradients of low and high waters between Planinsko polje and Ljubljance springs.




Fig. 5. Hydrogeologic units in the central part of Ljubljana Karst river basin. 1 — limestone, 2 — impermeable rocks (schists, etc.), 3 — dolomite, 4 — sediments on the karst paljes and moor, 5 — location of rupture survey, 6 — rupture diagrams, simplified, 7 — Idrija wrench fault (7) and other faults, 8 — hydrogeologic units: (1) — Vrhniko-Cerknica block, (2) — Rakov-Cerknica imbricate structure, (3) — Javorniki-Snežnik block, (4) — Hrušica block, (5) — Idrija-Ziri nappes area (all after S. BUSER 1965).

Sl. 6. Hidrogeološke enote v osrednjem delu kraškega porečja Ljubljane. 1 — apnenec, 2 — nepropustne kamnine (skrilavci in drugo), 3 — dolomit, 4 — naplavine na kraških poljih in barju, 5 — lokacija merjenih ruptur, 6 — diagrami ruptur, poenostavljeni, 7 — Idrijski zmik (7) in drugi prelomi, 8 — hidrogeološke enote: (1) — vrhniško-cerkniška gruda, (2) — rakovsko-cerkniška gruda, (3) — javorniško-snežniška gruda, (4) — hrušica, (5) — idrijsko-žirovsko ozetlje pokrovo (vse po S. BUSERJU 1965).





Fig. 1. Razpored izvornih Ljubljance, Lublje in Bistve
 Fig. 1. Ljubljana, Lubija and Bistva springs distribution



Andrej KRANJC: Skica ka referatu »Istorijske metode istraživanja kraških voda na Dolenjskem (Slovenija)«

Andrej KRANJC: A drawing enclosed to the report »Historical Methods of Water Investigations in Dolenjsko (Slovenia)«

LEGENDA:

1 = erozijski relief, 2 = kraško polje, 3 = površinski vodotok, 4 = kraško vrelo, 5 = izvorna pečina, 6 = ponor, 7 = ponorna pečina, 8 = podzemna vodena veza

Izvirne pečine: 1 — Podpeška pečina, 2 — Kumpoljska pečina, 3 — Potiskavška pečina, 4 — Lučka pečina, 5 — Krška pečina,

Ponorne pečine: 6 — Ponor Rašice, 7 — Vratnica, 8 — Finkova pečina, 9 — Tentera, 10 — Griška pečina, 11 — Zatočna pečina.

LEGEND:

1 = erosional relief, 2 = karst polje, 3 = surface water course, 4 = karst spring, 5 = spring cave, 6 = swallow hole (ponor), 7 = ponor cave, 8 = underground water connection,

Spring caves: 1 — Podpeška pečina, 2 — Kumpoljska pečina, 3 — Potiskavška pečina, 4 — Lučka pečina, 5 — Krška pečina,

Ponor caves: 6 — ponor Rašice, 7 — Vratnica, 8 — Finkova pečina, 9 — Tentera, 10 — Griška pečina, 11 — Zatočna pečina.

Fig 29





Fig. 31





1 - 4 = požiralniki

5 - 6 = bruhalnika

7 - 9 = jame

Fig. 32






Risba 11. Primeri presekov aktivnih in fosilnih izvirov jam

Drawing 11. Sections of active and fossil spring caves


Fig. 34




Riba 4. Potočna jama v NW koncu Ribniškega polja

Trondra 4. Potočna jama v NW koncu Ribniškega polja

Fig. 35





Sl. 27 Ponorī Rinžē, A - ponori pri Līvoldu, I - ošarvanti potīrātnīk, B - cēlotni tok Rinžē


Fig. 27 Rinža ponors, A - ponors near Līvold, I - dyed swallow holes, B - entire Rinža flow

Fig. 37


Figure 2. Hydrogeological sketch of the area between Kerey and Koyun


- Q - talus deposits, composed of fragments
- ls - brown mass, also fragments
- PIQ - siltstone (fine grained), also fragments - non-porous
- PI₂ - siltstone (fine grained), also fragments, non-porous
- E - siltstone (fine grained), also fragments, non-porous
- K - siltstone (fine grained), also fragments, non-porous
- ag - siltstone (fine grained), also fragments, non-porous
- J - siltstone (fine grained), also fragments, non-porous
- T - siltstone (fine grained), also fragments, non-porous
- F - siltstone (fine grained), also fragments, non-porous
- 1 - siltstone (fine grained), also fragments, non-porous
- 2 - siltstone (fine grained), also fragments, non-porous
- 3 - siltstone (fine grained), also fragments, non-porous
- 4 - siltstone (fine grained), also fragments, non-porous

- Q - talus deposits, permeable to impermeable
- ls - brown mass, permeable to impermeable
- PIQ - siltstone (fine grained), permeable to impermeable
- PI₂ - siltstone (fine grained), permeable to impermeable
- E - siltstone (fine grained), permeable to impermeable
- K - siltstone (fine grained), permeable to impermeable
- ag - siltstone (fine grained), permeable to impermeable
- J - siltstone (fine grained), permeable to impermeable
- T - siltstone (fine grained), permeable to impermeable
- F - siltstone (fine grained), permeable to impermeable
- 1 - siltstone (fine grained), permeable to impermeable
- 2 - siltstone (fine grained), permeable to impermeable
- 3 - siltstone (fine grained), permeable to impermeable
- 4 - siltstone (fine grained), permeable to impermeable




On OGA, 1:100,000 scale, Generalized in
Dance, 1980, 1:100,000 scale





Sl. 6. Vodna jama pri Cvišlerjih



Sl. 7. Vodna jama pri Ključni vasi